

Supplement 1. References for the studies on rehabilitation in older adults with hip fracture and cognitive impairment included in the reviewed literature

- S52. Watne LO, Torbergsen AC, Conroy S, Engedal K, Frihagen F, Hjorthaug GA, Juliebo V, Raeder J, Saltvedt I, Skovlund E, Wyller TB. The effect of a pre- and postoperative orthogeriatric service on cognitive function in patients with hip fracture: randomized controlled trial (Oslo Orthogeriatric Trial). *BMC Med* 2014;12:63. <https://doi.org/10.1186/1741-7015-12-63>
- S53. Huusko TM, Karppi P, Avikainen V, Kautiainen H, Sulkava R. Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: subgroup analysis of patients with dementia. *BMJ* 2000;321:1107-1111. <https://doi.org/10.1136/bmj.321.7269.1107>
- S54. Naglie G, Tansey C, Kirkland JL, Ogilvie-Harris DJ, Detsky AS, Etchells E, Tomlinson G, O'Rourke K, Goldlist B. Interdisciplinary inpatient care for elderly people with hip fracture: a randomized controlled trial. *CMAJ* 2002;167:25-32.
- S55. Jones GR, Miller TA, Petrella RJ. Evaluation of rehabilitation outcomes in older patients with hip fractures. *Am J Phys Med Rehabil* 2002;81:489-497. <https://doi.org/10.1097/00002060-200207000-00004>
- S56. McGilton KS, Mahomed N, Davis AM, Flannery J, Calabrese S. Outcomes for older adults in an inpatient rehabilitation facility following hip fracture (HF) surgery. *Arch Gerontol Geriatr* 2009;49:e23-e31. <https://doi.org/10.1016/j.archger.2008.07.012>
- S57. McGilton KS, Davis AM, Naglie G, Mahomed N, Flannery J, Jaglal S, Cott C, Stewart S. Evaluation of patient-centered rehabilitation model targeting older persons with a hip fracture, including those with cognitive impairment. *BMC Geriatr* 2013;13:136. <https://doi.org/10.1186/1471-2318-13-136>
- S58. Moseley AM, Sherrington C, Lord SR, Barracough E, St George RJ, Cameron ID. Mobility training after hip fracture: a randomised controlled trial. *Age Ageing* 2009;38:74-80. <https://doi.org/10.1093/ageing/afn217>
- S59. Rosler A, von Renteln-Kruse W, Muhlhan C, Frilling B. Treatment of dementia patients with fracture of the proximal femur in a specialized geriatric care unit compared to conventional geriatric care. *Z Gerontol Geriatr* 2012;45:400-403. <https://doi.org/10.1007/s00391-012-0299-1>
- S60. Stenvall M, Berggren M, Lundström M, Gustafson Y, Olofsson B. A multidisciplinary intervention program improved the outcome after hip fracture for people with dementia: subgroup analyses of a randomized controlled trial. *Arch Gerontol Geriatr* 2012;54:e284-e289. <https://doi.org/10.1016/j.archger.2011.08.013>
- S61. Giusti A, Barone A, Pioli G. Rehabilitation after hip fracture in patients with dementia. *J Am Geriatr Soc* 2007;55:1309-1310. <https://doi.org/10.1111/j.1532-5415.2007.01258.x>
- S62. Goldstein FC, Strasser DC. Hip fractures in older adults: cognitive factors related to a positive rehabilitation outcome. Emory University School, Department of Neurology and Rehabilitation Medicine; 1997.

- S63. Heruti RJ, Lusky A, Barell V, Ohry A, Adunsky A. Cognitive status at admission: does it affect the rehabilitation outcome of elderly patients with hip fracture? *Arch Phys Med Rehabil* 1999;80:432-436. [https://doi.org/10.1016/s0003-9993\(99\)90281-2](https://doi.org/10.1016/s0003-9993(99)90281-2)
- S64. Frances Horgan N, J Cunningham C. Impact of cognitive impairment on hip fracture outcome in older people. *Br J Ther Rehabil* 2003;10:228-232. <https://doi.org/10.12968/bjtr.2003.10.5.13547>
- S65. Lenze EJ, Skidmore ER, Dew MA, Butters MA, Rogers JC, Begley A, Reynolds CF, Munin MC. Does depression, apathy or cognitive impairment reduce the benefit of inpatient rehabilitation facilities for elderly hip fracture patients? *Gen Hosp Psychiatry* 2007;29:141-146. <https://doi.org/10.1016/j.genhosppsy.2007.01.001>
- S66. Penrod JD, Boockvar KS, Litke A, Magaziner J, Hannan EL, Halm EA, Silberzweig SB, Sean Morrison R, Orosz GM, Koval KJ, Siu AL. Physical therapy and mobility 2 and 6 months after hip fracture. *J Am Geriatr Soc* 2004;52:1114-1120. <https://doi.org/10.1111/j.1532-5415.2004.52309.x>
- S67. Rolland Y, Pillard F, Lauwers-Cances V, Busquere F, Vellas B, Lafont C. Rehabilitation outcome of elderly patients with hip fracture and cognitive impairment. *Disabil Rehabil* 2004;26:425-431. <https://doi.org/10.1080/09638280410001663148>
- S68. Uy C, Kurrle SE, Cameron ID. Inpatient multidisciplinary rehabilitation after hip fracture for residents of nursing homes: a randomised trial. *Australas J Ageing* 2008;27:43-44. <https://doi.org/10.1111/j.1741-6612.2007.00277.x>
- S69. Vidan M, Serra JA, Moreno C, Riquelme G, Ortiz J. Efficacy of a comprehensive geriatric intervention in older patients hospitalized for hip fracture: a randomized, controlled trial. *J Am Geriatr Soc* 2005;53:1476-1482. <https://doi.org/10.1111/j.1532-5415.2005.53466.x>
- S70. Al-Ani AN, Flodin L, Soderqvist A, Ackermann P, Samnegard E, Dalen N, Saaf M, Cederholm T, Hedstrom M. Does rehabilitation matter in patients with femoral neck fracture and cognitive impairment?: a prospective study of 246 patients. *Arch Phys Med Rehabil* 2010;91:51-57. <https://doi.org/10.1016/j.apmr.2009.09.005>
- S71. Chammout G, Kelly-Pettersson P, Hedbeck CJ, Boden H, Stark A, Mukka S, Skoldenberg O. Primary hemiarthroplasty for the elderly patient with cognitive dysfunction and a displaced femoral neck fracture: a prospective, observational cohort study. *Aging Clin Exp Res* 2021;33:1275-1283. <https://doi.org/10.1007/s40520-020-01651-8>
- S72. Karlsson A, Berggren M, Olofsson B, Stenvall M, Gustafson Y, Nordstrom P, Lindelof N. Geriatric interdisciplinary home rehabilitation after hip fracture in people with dementia: a subgroup analysis of a randomized controlled trial. *Clin Interv Aging* 2020;15:1575-1586. <https://doi.org/10.2147/CIA.S250809>
- S73. Uda K, Matsui H, Fushimi K, Yasunaga H. Intensive in-hospital rehabilitation after hip fracture surgery and activities of daily living in patients with dementia: retrospective analysis of a nationwide inpatient database. *Arch Phys Med Rehabil* 2019;100:2301-2307. <https://doi.org/10.1016/j.apmr.2019.06.019>

- S74. Sirbu PD, Forna N, Popescu DC, Tudor R, Friedl W, Ciubara A, Ciuntu BM, Berea G. Postoperative weight-bearing after uncemented bipolar hemiarthroplasty for femoral neck fractures in geriatric patients with dementia. *Broad Res Artif Intell Neurosci* 2019;10:66-71. <https://doi.org/10.70594/brain/v10.s1/9>
- S75. Raivio M, Korkala O, Pitkala K, Tilvis R. Rehabilitation outcome in hip-fracture: impact of weight-bearing restriction: a preliminary investigation. *Phys Occup Ther Geriatr* 2005;22:1-9. https://doi.org/10.1080/J148v22n04_01
- S76. Seitz DP, Gill SS, Austin PC, Bell CM, Anderson GM, Gruneir A, Rochon PA. Rehabilitation of older adults with dementia after hip fracture. *J Am Geriatr Soc* 2016;64:47-54. <https://doi.org/10.1111/jgs.13881>
- S77. Shyu YI, Tseng MY, Liang J, Tsai WC, Wu CC, Cheng HS. Interdisciplinary intervention decreases cognitive impairment for older Taiwanese with hip fracture: 2-year follow-up. *Int J Geriatr Psychiatry* 2013;28:1222-1231. <https://doi.org/10.1002/gps.3945>
- S78. Tseng MY, Yang CT, Liang J, Huang HL, Kuo LM, Wu CC, Cheng HS, Chen CY, Hsu YH, Lee PC, Shyu YL. A family care model for older persons with hip-fracture and cognitive impairment: a randomized controlled trial. *Int J Nurs Stud* 2021;120:103995. <https://doi.org/10.1016/j.ijnurstu.2021.103995>
- S79. Freter S, Koller K, Dunbar M, MacKnight C, Rockwood K. Translating delirium prevention strategies for elderly adults with hip fracture into routine clinical care: a pragmatic clinical trial. *J Am Geriatr Soc* 2017;65:567-573. <https://doi.org/10.1111/jgs.14568>
- S80. Marcantonio ER, Flacker JM, Wright RJ, Resnick NM. Reducing delirium after hip fracture: a randomized trial. *J Am Geriatr Soc* 2001;49:516-522. <https://doi.org/10.1046/j.1532-5415.2001.49108.x>
- S81. Bellelli G, Guerini F, Trabucchi M. Body weight-supported treadmill in the physical rehabilitation of severely demented subjects after hip fracture: a case report. *J Am Geriatr Soc* 2006;54:717-718. https://doi.org/10.1111/j.1532-5415.2006.00668_6.x